


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

BCR Polyclonal Antibody

Catalog No.E-AB-40429ReactivityH,M,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHCIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Images



Immunohistochemistry of paraffinembedded Human breast cancer using BCR Polycloanl Antibody at dilution of 1:200



Immunohistochemistry of paraffinembedded Mouse brain using BCR Polycloanl Antibody at dilution of 1:200



Immunohistochemistry of paraffinembedded Rat brain using BCR Polycloanl Antibody at dilution of 1:200

Immunogen Information

Immunogen Recombinant Human Breakpoint cluster region

protein

GeneID 613 **Swissprot** P11274

Synonyms ALL, bcr, BCR1, Breakpoint cluster region, CML,

PHL

Product Information

Buffer PBS with 0.05% Proclin300, 50% glycerol, pH7.3.

Purify Antigen Affinity Purification

Dilution IHC 1:100-1:300

Background

A reciprocal translocation between chromosomes 22 and 9 produces the Philadelphia chromosome, which is often found in patients with chronic myelogenous leukemia. The chromosome 22 breakpoint for this translocation is located within the BCR gene. The translocation produces a fusion protein which is encoded by sequence from both BCR and ABL, the gene at the chromosome 9 breakpoint. Although the BCR-ABL fusion protein has been extensively studied, the function of the normal BCR gene product is not clear. The protein has serine/threonine kinase activity and is a GTPase-activating protein for p21rac. Two transcript variants encoding different isoforms have been found for this gene.

For Research Use Only

Thank you for your recent purchase

If you would like to learn more about antibodies, please visit www.elabscience.com.

Focus on your research Service for life science