


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **BRCA1 Polyclonal Antibody**

Catalog No.E-AB-60016ReactivityHStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHCIsotypeIgG

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Images**



Immunohistochemistry of paraffinembedded human breast cancer using BRCA1 Polyclonal Antibody at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.

## **Immunogen Information**

**Immunogen** A synthetic peptide of human BRCA1

**GeneID** 672 **Swissprot** P38398

Synonyms BRCA1,BRCAI,BRCC1,BROVCA1,FANCS,IRIS,P

NCA4,PPP1R53,PSCP,RNF53

#### **Product Information**

Calculated MW 7kDa/78-85kDa/202-210kDa

**Observed MW** Refer to figures

**Buffer** PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Purify** Affinity purification **Dilution** IHC 1:50-1:200

## **Background**

This gene encodes a nuclear phosphoprotein that plays a role in maintaining genomic stability, and it also acts as a tumor suppressor. The encoded protein combines with other tumor suppressors, DNA damage sensors, and signal transducers to form a large multi-subunit protein complex known as the BRCA1-associated genome surveillance complex (BASC). This gene product associates with RNA polymerase II, and through the C-terminal domain, also interacts with histone deacetylase complexes. This protein thus plays a role in transcription, DNA repair of double-stranded breaks, and recombination. Mutations in this gene are responsible for approximately 40% of inherited breast cancers and more than 80% of inherited breast and ovarian cancers. Alternative splicing plays a role in modulating the subcellular localization and physiological function of this gene. Many alternatively spliced transcript variants, some of which are disease-associated mutations, have been described for this gene, but the full-length natures of only some of these variants has been described. A related pseudogene, which is also located on chromosome 17, has been identified.