


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# Cyclin D1 Polyclonal Antibody

Catalog No. E-AB-63508 Reactivity H,M,R Storage Store at -20°C. Avoid freeze / thaw cycles. Host Rabbit **Applications Isotype IgG** 

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

# **Images**



Western blot analysis of extracts of HCT116 cells using Cyclin D1 Polyclonal Antibody at dilution of 1:1000. HCT116 cells treated with different concentRations of RSPO1.

# **Immunogen Information**

**Immunogen** A synthetic peptide of human Cyclin D1

(NP\_444284.1).

GeneID 595 **Swissprot** P24385

**Synonyms** BCL1,D11S287E,PRAD1,U21B31,CCND1,Cyclin

D1,cyclin D1

#### **Product Information**

Calculated MW 33kDa **Observed MW** 37kDa

**Buffer** PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Purify** Affinity purification Dilution WB 1:500-1:2000

# **Background**

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance throughout the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with tumor suppressor protein Rb and the expression of this gene is regulated positively by Rb. Mutations, amplification and overexpression of this gene, which alters cell cycle progression, are observed frequently in a variety of tumors and may contribute to tumorigenesis.