


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **MYC Polyclonal Antibody**

Catalog No.E-AB-63525ReactivityM,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHCIsotypeIgG

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Images**



Immunohistochemistry of paraffinembedded Mouse liver using MYC Polyclonal Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffinembedded Rat heart using MYC Polyclonal Antibody at dilution of 1:100 (40x lens).

### **Immunogen Information**

**Immunogen** A synthetic peptide of human MYC (NP\_002458.2).

**GeneID** 4609 **Swissprot** P01106

**Synonyms** MRTL,MYCC,bHLHe39,c-Myc,MYC

#### **Product Information**

**Buffer** PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Purify** Affinity purification **Dilution** IHC 1:50-1:100

#### **Background**

The protein encoded by this gene is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. There is evidence to show that alternative translation initiations from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site result in the production of two isoforms with distinct N-termini. The synthesis of non-AUG initiated protein is suppressed in Burkitt's lymphomas, suggesting its importance in the normal function of this gene.