


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

TCF4 Polyclonal Antibody


Catalog No. E-AB-60206 Reactivity H,M,R Storage Store at -20°C. Avoid freeze / thaw cycles. Host Rabbit **Applications** WB.IF **Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of extracts of various cell lines using TCF4 Polyclonal Antibody at dilution of 1:1000.



Immunofluorescence analysis of C6 cells using TCF4 Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.

Immunogen Information

Immunogen Recombinant fusion protein of human TCF4

(NP 001077431.1).

GeneID 6925 **Swissprot** P15884

Synonyms TCF4,E2-2,FECD3,ITF-2,ITF2,PTHS,SEF-2,SEF2,S

EF2-1,SEF2-1A,SEF2-1B,SEF2-1D,TCF-4

Product Information

Calculated MW 48-71kDa **Observed MW** 60kDa

Buffer PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purify Affinity purification

Dilution WB 1:500-1:2000 IF 1:50-1:200

Background

This gene encodes transcription factor 4, a basic helix-loop-helix transcription factor. The encoded protein recognizes an Ephrussi-box ('Ebox') binding site ('CANNTG') - a motif first identified in immunoglobulin enhancers. This gene is broadly expressed, and may play an important role in nervous system development. Defects in this gene are a cause of Pitt-Hopkins syndrome. In addition, an intronic CTG repeat normally numbering 10-37 repeat units can expand to >50 repeat units and cause Fuchs endothelial corneal dystrophy. Multiple alternatively spliced transcript variants that encode different proteins have been described.