Elabscience®


C1 Inactivator Polyclonal Antibody

Catalog No.	E-AB-64154
Storage	Store at -20°C. Avoid freeze / thaw cycles.
Applications	WB,IHC


ReactivityH,M,RHostRabbitIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Images



Western blot analysis of extracts of Mouse heart using C1 Inactivator Polyclonal Antibody at dilution of 1:1000.



Immunohistochemistry of paraffinembedded Rat kidney using C1 Inactivator Polyclonal Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffinembedded Human breast cancer using C1 Inactivator Polyclonal Antibody at dilution of 1:100 (40x lens).

Immunogen Information

Immunogen	Recombinant fusion protein of human C1 Inactivator (NP_001027466.1).
GeneID	710
Swissprot	P05155
Synonyms	SERPING1,C1IN,C1INH,C1NH,HAE1,HAE2

Product Information

Calculated MW	49kDa/55kDa
Observed MW	60kDa
Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Purify	Affinity purification
Dilution	WB 1:500-1:2000 IHC 1:50-1:200

Background

This gene encodes a highly glycosylated plasma protein involved in the regulation of the complement cascade. Its protein inhibits activated C1r and C1s of the first complement component and thus regulates complement activation. Deficiency of this protein is associated with hereditary angioneurotic oedema (HANE). Alternative splicing results in multiple transcript variants encoding the same isoform.

For Research Use Only Thank you for your recent purchase

If you would like to learn more about antibodies,please visit www.elabscience.com.

Focus on your research Service for life science

Applications:WB-Western Blot IHC-Immunohistochemistry IF-Immunofluorescence IP-Immunoprecipitation FC-Flow cytometry ChIP-Chromatin Immunoprecipitation Reactivity: H-Human R-Rat M-Mouse Mk-Monkey Dg-Dog Ch-Chicken Hm-Hamster Rb-Rabbit Sh-Sheep Pg-Pig Z-Zebrafish X-Xenopus C-Cow.