


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

PSMD13 Polyclonal Antibody


Catalog No. E-AB-64203 Reactivity H,M,R Storage Store at -20°C. Avoid freeze / thaw cycles. Rabbit Host **Applications** WB.IF **Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of extracts of various cell lines using PSMD13 Polyclonal Antibody at 1:1000 dilution.



Immunofluorescence analysis of U2OS cells using PSMD13 Polyclonal Antibody Blue: DAPI for nuclear staining.

Immunogen Information

Immunogen Recombinant fusion protein of human PSMD13

5719 GeneID **Swissprot** O9UNM6

Synonyms PSMD13,HSPC027,Rpn9,S11,p40.5

Product Information

Calculated MW 42kDa **Observed MW** 43kDa

Buffer PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Purify Affinity purification

Dilution WB 1:500-1:2000,IF 1:50-1:200

Background

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a non-ATPase subunit of the 19S regulator. Two transcripts encoding different isoforms have been described.