


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

PSMC3 Polyclonal Antibody


Catalog No. E-AB-64570 Reactivity Η Store at -20°C. Avoid freeze / thaw cycles. Rabbit **Storage** Host **Applications** WB.IF **Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of extracts of various cell lines using PSMC3 Polyclonal Antibody at dilution of 1:1000.



Immunofluorescence analysis of U2OS cells using PSMC3 Polyclonal Antibody

Immunogen Information

Immunogen Recombinant fusion protein of human PSMC3

(NP 002799.3).

GeneID 5702 **Swissprot** P17980

Synonyms PSMC3,TBP1,ATPase 3

Product Information

Calculated MW 49kDa **Observed MW** 50kDa

Buffer PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purify Affinity purification

Dilution WB 1:500-1:2000 IF 1:50-1:200

Background

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases that have chaperone-like activity. This subunit may compete with PSMC2 for binding to the HIV tat protein to regulate the interaction between the viral protein and the transcription complex. A pseudogene has been identified on chromosome 9.