


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **PSMA6 Polyclonal Antibody**


Catalog No. E-AB-64698 Reactivity H,M,R Store at -20°C. Avoid freeze / thaw cycles. **Storage** Host Rabbit **Applications** WB.IF **Isotype IgG** 

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

# **Images**



Western blot analysis of extracts of various cell lines using PSMA6 Polyclonal Antibody at 1:1000 dilution.



Immunofluorescence analysis of MCF-7 cells using PSMA6 Polyclonal Antibody Blue: DAPI for nuclear staining.

# **Immunogen Information**

**Immunogen** Recombinant fusion protein of human PSMA6

GeneID 5687 **Swissprot** P60900

**Synonyms** PSMA6,IOTA,PROS27,p27K

#### **Product Information**

Calculated MW 18kDa/25kDa/27kDa

**Observed MW** 27kDa

**Buffer** PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Purify** Affinity purification

Dilution WB 1:500-1:2000,IF 1:50-1:200

# **Background**

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Multiple transcript variants encoding several different isoforms have been found for this gene. A pseudogene has been identified on the Y chromosome.