


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

NR3C1 Polyclonal Antibody


Reactivity Catalog No. E-AB-64779 Η Storage Store at -20°C. Avoid freeze / thaw cycles. Rabbit Host **Applications** WB.IF **Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of extracts of various cell lines using NR3C1 Polyclonal Antibody at dilution of 1:1000.



Immunofluorescence analysis of U2OS cells using NR3C1 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

Immunogen Information

Immunogen Recombinant fusion protein of human NR3C1

(NP 001191194.1).

GeneID 2908 **Swissprot** P04150

GCCR,GCR,GCRST,GR,GRL,NR3C1 **Synonyms**

Product Information

Calculated MW 49-85kDa **Observed MW** 105kDa

Buffer PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purify Affinity purification

Dilution WB 1:500-1:2000 IF 1:50-1:200

Background

This gene encodes glucocorticoid receptor, which can function both as a transcription factor that binds to glucocorticoid response elements in the promoters of glucocorticoid responsive genes to activate their transcription, and as a regulator of other transcription factors. This receptor is typically found in the cytoplasm, but upon ligand binding, is transported into the nucleus. It is involved in inflammatory responses, cellular proliferation, and differentiation in target tissues. Mutations in this gene are associated with generalized glucocorticoid resistance. Alternative splicing of this gene results in transcript variants encoding either the same or different isoforms. Additional isoforms resulting from the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-tonucleus trafficking patterns and distinct transcriptional activities (PMID:15866175).