


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **ATP6V1G3 Polyclonal Antibody**


Catalog No.E-AB-65007ReactivityMStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,IHCIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

# **Images**



Western blot analysis of extracts of Mouse kidney using ATP6V1G3 Polyclonal Antibody at dilution of 1:1000.



Immunohistochemistry of paraffinembedded Mouse kidney using ATP6V1G3 Polyclonal Antibody at dilution of 1:100 (40x lens).

# **Immunogen Information**

**Immunogen** Recombinant fusion protein of human ATP6V1G3

(NP\_573569.1).

**GeneID** 127124 **Swissprot** Q96LB4

**Synonyms** ATP6V1G3,ATP6G3,Vma10

#### **Product Information**

Calculated MW 6kDa/13kDa/14kDa

**Observed MW** 14kDa

**Buffer** PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Purify** Affinity purification

**Dilution** WB 1:500-1:2000 IHC 1:50-1:100

# **Background**

This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c" and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This gene encodes one of three G subunit proteins. Transcript variants encoding different isoforms have been found for this gene.