


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

ATP6V1A Polyclonal Antibody

Catalog No. E-AB-65078 Reactivity M Store at -20°C. Avoid freeze / thaw cycles. Rabbit **Storage** Host **Applications Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of extracts of Mouse brain using ATP6V1A Polyclonal Antibody at dilution of 1:1000.

Immunogen Information

Immunogen A synthetic peptide of human ATP6V1A

(NP_001681.2).

GeneID 523 **Swissprot** P38606

Synonyms ATP6V1A,ATP6A1,ATP6V1A1,HO68,VA68,VPP2,

Vma1,ARCL2D

Product Information

Calculated MW 64kDa/68kDa

Observed MW 74kDa

Buffer PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purify Affinity purification Dilution WB 1:500-1:2000

Background

This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c', and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This encoded protein is one of two V1 domain A subunit isoforms and is found in all tissues. Transcript variants derived from alternative polyadenylation exist.