


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

PSMC5 Polyclonal Antibody


Catalog No.E-AB-60331ReactivityH,M,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,IFIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of extracts of various cell lines using PSMC5 Polyclonal Antibody at dilution of 1:1000.



Immunofluorescence analysis of MCF-7 cells using PSMC5 Polyclonal Antibody

Immunogen Information

Immunogen Recombinant fusion protein of human PSMC5

(NP_002796.4).

GeneID 5705 **Swissprot** P62195

Synonyms PSMC5,S8,SUG-1,SUG1,TBP10,TRIP1,p45,p45/SU

G,ATPase 5

Product Information

Calculated MW 44kDa/45kDa

Observed MW 46kDa

Buffer PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purify Affinity purification

Dilution WB 1:500-1:2000 IF 1:50-1:200

Background

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases which have a chaperone-like activity. In addition to participation in proteasome functions, this subunit may participate in transcriptional regulation since it has been shown to interact with the thyroid hormone receptor and retinoid X receptor-alpha. Two transcript variants encoding different isoforms have been found for this gene.