


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

DDB1 Polyclonal Antibody

Catalog No. E-AB-65453 Reactivity H,M,R Store at -20°C. Avoid freeze / thaw cycles. **Storage** Host Rabbit **Applications Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of extracts of various cell lines using DDB1 Polyclonal Antibody at dilution of 1:3000.

Immunogen Information

A synthetic peptide of human DDB1 (NP 001914.3). **Immunogen**

GeneID 1642 **Swissprot** O16531

Synonyms DDB1.DDBA.UV-DDB1.XAP1.XPCE.XPE.XPE-BF

Product Information

Calculated MW 50kDa/126kDa

Observed MW 127kDa

Buffer PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purify Affinity purification Dilution WB 1:500-1:2000

Background

The protein encoded by this gene is the large subunit (p127) of the heterodimeric DNA damage-binding (DDB) complex while another protein (p48) forms the small subunit. This protein complex functions in nucleotide-excision repair and binds to DNA following UV damage. Defective activity of this complex causes the repair defect in patients with xeroderma pigmentosum complementation group E (XPE) - an autosomal recessive disorder characterized by photosensitivity and early onset of carcinomas. However, it remains for mutation analysis to demonstrate whether the defect in XPE patients is in this gene or the gene encoding the small subunit. In addition, Best vitelliform mascular dystrophy is mapped to the same region as this gene on 11q, but no sequence alternations of this gene are demonstrated in Best disease patients. The protein encoded by this gene also functions as an adaptor molecule for the cullin 4 (CUL4) ubiquitin E3 ligase complex by facilitating the binding of substrates to this complex and the ubiquitination of proteins.