


CBR4 Polyclonal Antibody

Catalog No.	E-AB-63190	Reactivity	H,R
Storage	Store at -20°C. Avoid freeze / thaw cycles.	Host	Rabbit
Applications	WB	Isotype	IgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of extracts of various cell lines using CBR4 Polyclonal Antibody at 1:1000 dilution.

Immunogen Information

Immunogen	Recombinant fusion protein of human CBR4
GeneID	84869
Swissprot	Q8N4T8
Synonyms	CBR4,SDR45C1

Product Information

Calculated MW	18kDa/25kDa
Observed MW	25kDa
Buffer	PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Purify	Affinity purification
Dilution	WB 1:200-1:2000

Background

Component of the heterotetramer complex KAR (3-ketoacyl-[acyl carrier protein] reductase or 3-ketoacyl-[ACP] reductase that forms part of the mitochondrial fatty acid synthase (mtFAS. Beta-subunit of the KAR heterotetramer complex, responsible for the 3-ketoacyl-ACP reductase activity of the mtFAS, reduces 3-oxoacyl-[ACP] to (3R-hydroxyacyl-[ACP] in a NADPH-dependent manner with no chain length preference, thereby participating in mitochondrial fatty acid biosynthesis. The homotetramer has NADPH-dependent quinone reductase activity (in vitro, hence could play a role in protection against cytotoxicity of exogenous quinones. As a heterotetramer, it can also reduce 9,10-phenanthrenequinone, 1,4-benzoquinone and various other o-quinones and p-quinones (in vitro).

For Research Use Only

Thank you for your recent purchase.
If you would like to learn more about antibodies, please visit www.elabscience.com.

Focus on your research Service for life science

Applications: WB-Western Blot IHC-Immunohistochemistry IF-Immunofluorescence IP-Immunoprecipitation FC-Flow cytometry ChIP-Chromatin Immunoprecipitation Reactivity: H-Human R-Rat M-Mouse Mk-Monkey Dg-Dog Ch-Chicken Hm-Hamster Rb-Rabbit Sh-Sheep Pg-Pig Z-Zebrafish X-Xenopus C-Cow.