


Tel:1-832-243-6086 Fax:1-832-243-6017 techsupport@elabscience.com Website: www.elabscience.com

PSMD9 Polyclonal Antibody

Catalog No. E-AB-60921 Reactivity H,M,R Store at -20°C. Avoid freeze / thaw cycles. **Storage** Host Rabbit **Applications Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of extracts of various cell lines using PSMD9 Polyclonal Antibody at 1:1000 dilution.

Immunogen Information

Immunogen Recombinant fusion protein of human PSMD9

GeneID 5715 O00233 **Swissprot**

Synonyms PSMD9,Rpn4,p27

Product Information

Calculated MW 13kDa/22kDa/24kDa

Observed MW 25kDa

Buffer PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Purify Affinity purification Dilution WB 1:500-1:2000

Background

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a non-ATPase subunit of the 19S regulator. Three transcript variants encoding two different isoforms have been found for this gene.