


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

HAO1 Polyclonal Antibody


Reactivity Catalog No. E-AB-61988 H,M,R Storage Store at -20°C. Avoid freeze / thaw cycles. Rabbit Host **Applications** WB,IHC **Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of extracts of various cell lines using HAO1 Polyclonal Antibody at dilution of 1:500.



Immunohistochemistry of paraffinembedded Human liver cancer using HAO1 Polyclonal Antibody at dilution of 1:200 (40x lens).

Immunogen Information

Immunogen Recombinant fusion protein of human HAO1

(NP_060015.1).

GeneID 54363 **Swissprot** Q9UJM8

HAO1,GOX,GOX1,HAOX1 **Synonyms**

Product Information

Calculated MW 40kDa **Observed MW** 41kDa

Buffer PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purify Affinity purification

Dilution WB 1:500-1:2000 IHC 1:50-1:200

Background

This gene is one of three related genes that have 2-hydroxyacid oxidase activity yet differ in encoded protein amino acid sequence, tissue expression and substrate preference. Subcellular location of the encoded protein is the peroxisome. Specifically, this gene is expressed primarily in liver and pancreas and the encoded protein is most active on glycolate, a two-carbon substrate. The protein is also active on 2-hydroxy fatty acids. The transcript detected at high levels in pancreas may represent an alternatively spliced form or the use of a multiple near-consensus upstream polyadenylation site.