


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **RPL14 Polyclonal Antibody**


Reactivity Catalog No. E-AB-61324 H,M,R Store at -20°C. Avoid freeze / thaw cycles. Rabbit **Storage** Host **Applications** WB.IF **Isotype IgG** 

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Images**



Western blot analysis of extracts of various cell lines using RPL14 Polyclonal Antibody at 1:1000 dilution.



Immunofluorescence analysis of MCF7 cells using RPL14 Polyclonal Antibody Blue: DAPI for nuclear staining.

## **Immunogen Information**

**Immunogen** Recombinant fusion protein of human RPL14

9045 GeneID P50914 **Swissprot** 

**Synonyms** RPL14.CAG-ISL-7.CTG-B33.L14.RL14.hRL14

#### **Product Information**

Calculated MW 23kDa **Observed MW** 26kDa

**Buffer** PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Purify** Affinity purification

Dilution WB 1:500-1:2000,IF 1:50-1:200

#### **Background**

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L14E family of ribosomal proteins. It contains a basic region-leucine zipper (bZIP)-like domain. The protein is located in the cytoplasm. This gene contains a trinucleotide (GCT) repeat tract whose length is highly polymorphic; these triplet repeats result in a stretch of alanine residues in the encoded protein. Transcript variants utilizing alternative polyA signals and alternative 5'-terminal exons exist but all encode the same protein. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.