


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

SETMAR Polyclonal Antibody

Catalog No.E-AB-61336ReactivityHStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHCIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Immunohistochemistry of paraffinembedded human colon carcinoma using SETMAR Polyclonal Antibody at dilution of 1:100 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry of paraffinembedded rat spleen using SETMAR Polyclonal Antibody at dilution of 1:100 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

Immunogen Information

Immunogen Recombinant fusion protein of human SETMAR

GeneID 6419 **Swissprot** Q53H47

Synonyms SETMAR, METNASE, Mar1

Product Information

Calculated MW 40kDa/62kDa/78kDa **Observed MW** Refer to Figures

Buffer PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Purify Affinity purification **Dilution** IHC 1:50-1:200

Background

This gene encodes a fusion protein that contains an N-terminal histonelysine N-methyltransferase domain and a C-terminal mariner transposase domain. The encoded protein binds DNA and functions in DNA repair activities including non-homologous end joining and double strand break repair. The SET domain portion of this protein specifically methylates histone H3 lysines 4 and 36. This gene exists as a fusion gene only in anthropoid primates, other organisms lack mariner transposase domain. Alternate splicing results in multiple transcript variants.