


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **CRYGS Polyclonal Antibody**

Catalog No.E-AB-62697ReactivityH,M,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWBIsotypeIgG

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

# **Images**



Western blot analysis of extracts of various cell lines using CRYGS Polyclonal Antibody at dilution of 1:4000.

# **Immunogen Information**

Immunogen Recombinant fusion protein of human CRYGS

(NP\_060011.1).

**GeneID** 1427 **Swissprot** P22914

Synonyms CRYGS,CRYG8,CTRCT20

#### **Product Information**

Calculated MW 21kDa Observed MW 21kDa

**Buffer** PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Purify** Affinity purification **Dilution** WB 1:500-1:2000

# **Background**

Crystallins are separated into two classes: taxon-specific, or enzyme, and ubiquitous. The latter class constitutes the major proteins of vertebrate eye lens and maintains the transparency and refractive index of the lens. Since lens central fiber cells lose their nuclei during development, these crystallins are made and then retained throughout life, making them extremely stable proteins. Mammalian lens crystallins are divided into alpha, beta, and gamma families; beta and gamma crystallins are also considered as a superfamily. Alpha and beta families are further divided into acidic and basic groups. Seven protein regions exist in crystallins: four homologous motifs, a connecting peptide, and N- and C-terminal extensions. Gamma-crystallins are a homogeneous group of highly symmetrical, monomeric proteins typically lacking connecting peptides and terminal extensions. They are differentially regulated after early development. This gene encodes a protein initially considered to be a betacrystallin but the encoded protein is monomeric and has greater sequence similarity to other gamma-crystallins. This gene encodes the most significant gamma-crystallin in adult eye lens tissue. Whether due to aging or mutations in specific genes, gamma-crystallins have been involved in cataract formation.

For Research Use Only

Thank you for your recent purchase

If you would like to learn more about antibodies, please visit www.elabscience.com.

Focus on your research Service for life science