


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

CCL2 Polyclonal Antibody


Catalog No.E-AB-70085ReactivityH,MStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHCIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Images



Immunohistochemistry analysis of paraffin-embedded human colon using CCL2 Polyclonal Antibody at dilution of 1:500.



Immunohistochemistry analysis of paraffin-embedded human lung cancer using CCL2 Polyclonal Antibody at dilution of 1:300.



Immunohistochemistry analysis of paraffin-embedded mouse liver using CCL2 Polyclonal Antibody at dilution of 1:300.

Immunogen Information

Immunogen Recombinant protein corresponding to Mouse MCP-1

Swissprot P13500,P10148

Synonyms CCL2, GDCF-2, HC11, HSMCR30, MCAF, MCP1,

SCYA2, SMC-CF, C-C motif chemokine ligand 2

Product Information

Buffer PBS with 0.02% sodium azide, 1% protective protein

and 50% glycerol, pH7.4

Purify Affinity purification
Dilution IHC 1:200-1:800

Background

This gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXC subfamily of cytokines. Members of this subfamily are characterized by two cysteines separated by a single amino acid. This cytokine displays chemotactic activity for monocytes and basophils but not for neutrophils or eosinophils. It has been implicated in the pathogenesis of diseases characterized by monocytic infiltrates, like psoriasis, rheumatoid arthritis and atherosclerosis. It binds to chemokine receptors CCR2 and CCR4.

For Research Use Only

Thank you for your recent purchase

If you would like to learn more about antibodies, please visit www.elabscience.com.

Focus on your research Service for life science