


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

BMP2 Polyclonal Antibody


Catalog No.E-AB-70111ReactivityH,M,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,IHCIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western Blot analysis of various samples using BMP2 Polyclonal Antibody at dilution of 1:1000.



Immunohistochemistry analysis of paraffin-embedded mouse bone marrow using BMP2 Polyclonal Antibody at dilution of 1:300.

Immunogen Information

Immunogen KLH conjugated Synthetic peptide corresponding to

Mouse BMP2

Swissprot P12643,P21274,P49001

Synonyms BMP2, BDA2, BMP2A, bone morphogenetic protein

2

Product Information

Calculated MW 45kDa **Observed MW** 45kDa

Buffer PBS with 0.02% sodium azide, 1% protective protein

and 50% glycerol, pH7.4

Purify Affinity purification

Dilution WB 1:500-1:2000, IHC 1:200-1:800

Background

Bone morphogenetic protein-2 (BMP-2) is a member of the transforming growth factor-beta (TGFB) superfamily. BMP2 is synthesized as a 60 kDa precursor that is processed in the secretory pathway to a small 18 kDa monomer; 2 monomers then associate to form the active 30 kDa homodimer, which binds to its receptor. There is also a 40-45 kDa form of BMP2, as an amino-terminal propeptide. BMP2 can induce bone formation and regeneration during early embryonic development. It is involved in the hedgehog pathway, TGF beta signaling pathway, and cytokine-cytokine receptor interaction.