


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **ROCK1 Polyclonal Antibody**


Catalog No.E-AB-70124ReactivityM,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHCIsotypeIgG

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

# **Images**



Immunohistochemistry analysis of paraffin-embedded mouse kidney using ROCK1 Polyclonal Antibody at dilution of 1:300.



Immunohistochemistry analysis of paraffin-embedded rat spleen using ROCK1 Polyclonal Antibody at dilution of 1:300.

# **Immunogen Information**

Immunogen KLH conjugated Synthetic peptide corresponding to

Mouse Rock1

**Swissprot** P70335,Q63644

**Synonyms** P160ROCK, ROCK-I, Rho associated coiled-coil

containing protein kinase 1

#### **Product Information**

**Buffer** PBS with 0.02% sodium azide, 1% protective protein

and 50% glycerol, pH7.4

Purify Affinity purification
Dilution IHC 1:100-1:400

# **Background**

This gene encodes a protein serine/threonine kinase that is activated when bound to the GTP-bound form of Rho. The small GTPase Rho regulates formation of focal adhesions and stress fibers of fibroblasts, as well as adhesion and aggregation of platelets and lymphocytes by shuttling between the inactive GDP-bound form and the active GTP-bound form. Rho is also essential in cytokinesis and plays a role in transcriptional activation by serum response factor. This protein, a downstream effector of Rho, phosphorylates and activates LIM kinase, which in turn, phosphorylates cofilin, inhibiting its actin-depolymerizing activity.